

**Almond Board of California Disease Forecasts 2024**  
in cooperation with the University of California and Semios

**Table 1.** 7-day disease risk forecasts for Mon., June 10 through Mon., June 17, 2024\*

No.	County	Region	Anthracoze (value, date, color code)^	Bacterial spot (value, date, color code)^	Alternaria leaf spot (value, date, color code)^	Almond scab sporulation level (date, T, LW value, Precip.)^
1	Butte	West	0	0	0; Seasonal-DI = 74; up 13	0
2	Colusa	East	0	0	0	0
3	Fresno	Central	0	0	0	0
4	Fresno	East	0	0	0; Seasonal-DI = 26; up 10	0
5	Fresno	West	0	0	0	0
6	Kern	Central	0	0	0	0
7	Kern	East	0	0	0; Seasonal-DI = 38; up 4	0
8	Kern	West	0	0, 7-DI of 4.27 to 2.14 from 6/10 to 6/14	0	0
9	Madera	Central	0	0	0; Seasonal-DI = 16; up 4	0
10	Merced	Central	0	0	0	0
11	Stanislaus	Central	0	0	0	0
12	Stanislaus	East	0	0	0	0
13	Stanislaus	West	0	0	0	0

\* - 7-day forecasts are based on temperature (inside- and outside-canopy measurements), precipitation, and leaf wetness which are powered by the Semios® precision farming platform. 5-, 7-, and 21-day disease indices (DI) are also shown that provide the previous risk for a region.

^ - Numerical risk is scaled as follows: 0 = no risk, 1 = action threshold (Note: values may exceed 1 due to hourly accumulations). Color code risk: yellow = low, orange = moderate, red = high.

### Industry Advisory - Summary for Selected Almond Growing Regions

No appreciable rainfall and moderate average temperatures occurred in all regions last week (Table 3). Average daily leaf wetness durations ranged from 10.4 h in Butte-W, 8.6 h in Fresno-E, 5.0 h in Madera-C, 3.7 h in Stanislaus-W, and 2.6 h in Kern-E, whereas  $\leq 1.6$  h occurred in other regions. This resulted in increases in DSV values for Alternaria leaf spot (ALS) of 13 in Butte-W, 10 in Fresno-E, and 4 in Kern-E and Madera-C, with seasonal DSVs reaching 74, 26, 38, and 16, respectively. Last week's forecast again called for a low risk, but high leaf wetness hours contributed to a high risk for Alternaria leaf spot in Butte-W, and Fresno-E. Other regions still have a 0 seasonal DSV, indicating low risk for ALS. High daily leaf wetness durations with moderate avg. temperatures ( $>11.0^\circ$  avg. lows to  $<41.9^\circ$ C avg. highs) resulted in a low forecasted risk for scab sporulation in Butte-W, Madera-C, and Fresno-E (Table 3). Higher leaf wetness, however, actually occurred than what was forecasted in Butte-W and Fresno-E, as well as Kern-E, Madera-C, and Stanislaus-W, which increased the risk not only for Alternaria leaf spot but also scab sporulation in these locations. Thus, fungicide applications for ALS and scab are warranted in Butte-W, Fresno-E, Madera-C, and Kern-E, if applications have not been already applied. ALS fungicide treatments are suggested at DSV intervals of 8 to 12 depending on the intended level of disease management. A fungicide treatment is persistent for three weeks under low rainfall conditions. Thus, a maximum of 3 applications will provide 9 weeks of protection. With no appreciable rainfall, the risk for anthracnose and bacterial spot risk was 0 for all regions except in Kern-W where bacterial spot risk reached a 7-DI high of 4.27 on 6-8-24.

For the coming week, no precipitation, low daily leaf wetness hours, moderate to high temperatures and moderate to low humidity are forecasted for all regions (Table 2). Daily leaf wetness values of  $\leq 0.3$  h are forecasted for all regions with 0.3 h for Colusa-E. Temperatures are holding (daily averages between  $22.8^\circ$  and  $31.7^\circ$ C) with daily maximum values of up to  $41.9^\circ$ C ( $107.4^\circ$ F). With no rainfall and low leaf wetness forecasted in the coming week, zero to low risk is predicted for anthracnose, bacterial spot, and almond scab sporulation as shown in Table 1. Alternaria leaf spot risks are also forecasted as low (0 values), but heavy dews or irrigation practices have increased the daily wetness periods in selected regions for the last several weeks. Thus, contrary to the forecast, it seems that the risk for Alternaria and scab is high for Butte-W and Fresno-E with extended leaf wetness periods occurring in each of the last several weeks including last week as shown in Tables 1 and 3 and continues to increase in Kern-E and Madera-C.

The website <https://www.ag-radar.com> (password: Almondboard2022) displays actual and forecasted disease risk assessments for each region. Because these are regional forecasts, actual and predicted precipitation may vary among locations within each region. Additionally, historical records and experience for specific locations should be considered. This advisory will be updated weekly. The website "2022 Fungicide Efficacy Tables" is available to optimize fungicide selection and applications (<http://ipm.ucanr.edu/PDF/PMG/fungicideefficacytiming.pdf>).

**Table 2.** Forecasted weather for Mon., June 10, through Mon., June 17, 2024\*

No.	County	Region	Avg Temp (in canopy) (Avg)°C	Avg Humidity (Avg) (%)	Total Precip. (mm)	Leaf Wetness (hours/day)
1	Butte	West	24.2 – 31.2 (26.9)	21.5 – 59.3 (32.7)	0	0
2	Colusa	East	25.0 – 30.8 (26.8)	22.0 – 45.4 (33.6)	0	0.3
3	Fresno	Central	23.9 – 29.5 (26.5)	27.2 – 53.1 (36.7)	0	0
4	Fresno	East	24.1 – 30.2 (27.0)	31.0 – 50.4 (39.4)	0	0
5	Fresno	West	22.8 – 28.1 (25.4)	21.4 – 44.3 (31.6)	0	0
6	Kern	Central	23.4 – 29.1 (26.2)	26.0 – 52.6 (34.0)	0	0
7	Kern	East	24.4 – 31.7 (27.8)	28.8 – 47.4 (34.4)	0	0
8	Kern	West	24.3 – 30.0 (26.9)	24.2 – 45.2 (33.1)	0	0
9	Madera	Central	23.3 – 29.0 (25.9)	30.9 – 53.6 (39.6)	0	0
10	Merced	Central	23.4 – 29.3 (25.6)	31.3 – 55.5 (39.4)	0	0
11	Stanislaus	Central	23.3 – 28.1 (25.0)	31.1 – 48.4 (37.7)	0	0
12	Stanislaus	East	22.9 – 28.4 (25.1)	34.5 – 56.5 (41.1)	0	0
13	Stanislaus	West	23.9 – 28.7 (25.5)	27.2 – 51.3 (35.4)	0	0


**Table 3.** Previous week’s actual weather for Mon., June 3, through Sun., June 9, 2024\*

No.	County	Region	Avg Temp (in canopy) (Avg)°C	Avg Humidity (Avg) (%)	Total Precip. (mm)	Leaf Wetness (hours/day)
1	Butte	West	23.2 – 28.0 (25.5)	55.8 – 68.1 (61.5)	0	10.4
2	Colusa	East	21.5 – 26.6 (24.2)	35.3 – 55.1 (45.1)	0	0.6
3	Fresno	Central	22.4 – 28.5 (24.9)	49.8 – 62.3 (57.5)	0	0
4	Fresno	East	22.7 – 29.4 (25.7)	47.7 – 58.1 (53.9)	0	8.6
5	Fresno	West	23.2 – 30.3 (26.4)	37.4 – 57.1 (47.5)	0	0
6	Kern	Central	22.8 – 28.8 (25.0)	53.0 – 63.7 (59.2)	0	0
7	Kern	East	23.2 – 29.1 (26.4)	47.8 – 60.6 (55.7)	0	2.6
8	Kern	West	23.1 – 29.4 (26.0)	44.2 – 58.4 (51.9)	0.6	0
9	Madera	Central	23.2 – 29.1 (25.2)	50.3 – 62.8 (56.9)	0	5.0
10	Merced	Central	21.8 – 27.1 (23.7)	57.6 – 65.5 (61.3)	0	1.6
11	Stanislaus	Central	21.4 – 28.4 (24.6)	38.9 – 56.9 (45.6)	0	0.4
12	Stanislaus	East	20.7 – 28.1 (24.0)	53.2 – 69.1 (57.7)	0	0
13	Stanislaus	West	21.1 – 27.6 (24.0)	45.2 – 57.7 (50.2)	0	3.7

**Note:** Data in these tables were generated using the RADAR on-line forecasted report powered by the Semios® precision farming platform.

Fig. 1. Maps of counties and regions.

Butte West



Colusa East



Fresno Central



Fresno East



Fresno West





Fig. 2. Maps of counties and regions.

Kern West



Kern East



Kern Central



Stanislaus Central



Stanislaus East



Stanislaus West

